



Compounding the complexity

The impact of comorbidities on the management of non-communicable diseases

Authors:

Percept Actuaries and Consultants

17 February 2026

17

02

26



Research briefs on non-communicable diseases in South Africa

We have developed a series of briefs aiming to explain, explore and quantify the burden of non-communicable diseases (NCDs) in South Africa. Throughout the briefs, we draw together both existing quantitative data as well as emerging qualitative data. The primary qualitative data presented in the form of vignettes, has been collected by Dr Beth Vale, through in-depth ethnographic research.

These briefs are timely, given the rising global burden, and particularly the rising burden in low- and middle-income countries (LMICs), of NCDs. Given South Africa's high prevalence of HIV, there has also been a recent focus on the link between HIV and NCDs as the population living with HIV grow increasingly older with the successful uptake of antiretrovirals. An ageing population is more at risk for NCDs as we will explain in our briefs. As we move towards universal health coverage (UHC) it is imperative that we understand the needs of our population now and how these may change going forward. We have produced twelve briefs in this series.

This work was supported by the following partners, although the views are the authors' own:

- Actuarial Society of South Africa (ASSA): ASSA has an interest in being part of the development of high-quality evidence to support resource allocation and decision-making and the interplay between the supply and demand sides of the health system.
- RGA Reinsurance Company of South Africa Ltd (RGA): RGA has an interest in the ways in which life insurance can be responsive to the changing burden of disease and the ways in which we can use data to drive decision-making.
- Board of Healthcare Funders (BHF): BHF has an interest in an affordable, accessible, and high-quality private health sector.

Take-home messages

- The prevalence of co- or multimorbid chronic diseases in South Africa is likely substantially underestimated.
- Many chronic illnesses are associated with one another in a multitude of ways. This makes disease management more complex for both clinicians and clients, which in turn negatively impacts quality of care.
- Limitations in publicly available data for the uninsured population makes estimating the prevalence of co- and multimorbidity challenging. However, survey data shows that approximately 11% of people have at least two chronic conditions. The prevalence of co- and multimorbidity is higher in women than in men, and three quarters of people with diabetes have a comorbidity.
- Hospitalisation is 2.73 times more likely in the previous six months for people with multimorbid chronic diseases.
- A person-centred approach which looks at treating the whole person rather than just the disease is needed to manage comorbidities, and this should be supported by more integrated chronic disease management clinical guidelines and tools to measure and manage polypharmacy.



Introduction

This brief focuses on comorbidity and multimorbidity of chronic conditions. For our purposes, we define comorbidity as an individual having two chronic conditions at the same time, and multimorbidity as an individual having more than two chronic conditions at the same time.¹ We use the term chronic conditions, rather than non-communicable diseases (NCDs) to include HIV which is a communicable disease but also chronic. We do not include medium-duration conditions like tuberculosis (TB) in our definition of comorbidity, as TB is curable. Therefore, in this brief, we focus on long term, chronic conditions that need to be managed using both medical, behavioural, and/or social interventions for the remainder of a person's life.

In previous briefs we explained how NCDs are a growing global concern and the leading cause of death worldwide in recent years.² However, an area that is still being researched is the impact of comorbidity on health outcomes, and the relationship between different types of chronic diseases and their risk factors.

Complex causality and compounded costs of care

Previous briefs in this series have explored the causes of chronic diseases. Biological factors like age and sex play a strong role in the onset and type of disease. Similarly, socio-economic status and living conditions also contribute to a person's susceptibility to chronic conditions. These same factors also play a role in comorbidity.

However, some chronic conditions are also widely accepted to be inherently linked due to the **crossover of the factors that first give rise to them (causative factors)**. Examples of these conditions are heart disease, hyperlipidaemia (high cholesterol) and hypertension.³ Therefore, if a person displays the risk factors linked to these conditions, they are also likely to experience all three conditions simultaneously.

Relationships also exist between **infectious diseases and NCDs**. For example, Antiretroviral Therapy (ART), used to treat HIV, has been linked to increased cardiovascular disease, and the Human Papillomavirus (HPV), a sexually transmitted infection (STI), can cause cervical cancer.⁴ Both these conditions are particularly relevant for South Africa, given the high prevalence of HIV and STIs. Therefore, a person's experience of infectious diseases can also contribute to their susceptibility for NCDs. This linkage goes in both directions. For example, diabetes makes people more susceptible to acquiring TB and pneumonia, both of which are infectious diseases.⁴

Certain combinations of chronic conditions exhibit what is referred to as superadditive costs of treatment. Superadditivity is when the sum of two elements leads to a value greater than just the additive sum of the two elements. In this sense having two chronic diseases does not simply sum to the cost of treating these two conditions.⁵ For example, if a patient has both diabetes and heart disease, and it costs R500 to treat diabetes and R500 to treat heart disease, the cost of treating both conditions together will be greater than R1,000. However, not all chronic conditions are super-additive. Finding the ones that are and targeting prevention efforts at these can be substantially more effective than a single disease prevention lens.⁵

One study found that in ten selected chronic conditions^a (which resulted in 45 combinations of comorbidities which were analysed), 41 combinations of these chronic conditions were found to be superadditive.⁵ Diabetes was found to be a particularly significant driver of superadditive costs when co-occurring with another chronic condition; it was found to significantly increase the aggregate cost of treating chronic kidney disease, heart disease, respiratory illnesses, and stroke.⁵ A study of health system

^a This study specifically looked at stroke, heart disease, cancers (breast, lung, colorectal, stomach, liver, kidney, pancreatic, and oesophageal), diabetes, chronic kidney disease, respiratory illness (chronic obstructive pulmonary disease and asthma), cirrhosis, major depression, neurological disorders, and alcohol use disorders.⁵ Combinations of comorbidities included stroke, heart disease, cancers, diabetes, and chronic kidney disease for each of the ten chronic conditions, respectively.⁵




costs for individual and comorbid NCDs in New Zealand found that 23.8% of total health expenditure on NCDs could be attributed to the superadditive nature of certain combinations of comorbid chronic conditions.⁶

Cortaredona and Ventelou (2017) argue that, “When comorbidity exists and where super-additivity is involved, a given preventive policy leads to greater monetary savings than the costs associated with the single diagnosis, meaning that, **the returns from the action are generally underestimated.**”⁵ Therefore, we need to understand both the relationship between risk factors for different conditions and the impact on health needs when someone lives with more than one chronic condition. When both these factors are considered, it becomes clear that our prevention efforts and service delivery modes need to adapt to urgently stem the tide of, and better support individuals with, comorbidities.

Comorbidities make care more complicated

Figure 1 below shows how comorbidity influences the complexity of care a client would require from the health system, making quality care more difficult to render.³ It shows that when there are a number of chronic conditions present at the same time in one patient, care becomes more complicated due to the specific characteristics of these conditions independently, and also how these conditions interact with one another.³

Figure 1: Conceptual framework of the influence of comorbidities on clinical complexity and quality of care for patients³



When certain chronic conditions are more dominant in terms of the intensity of the symptoms or the onset of one chronic condition is related to the onset of another, it makes diagnosis and management of these conditions more challenging from a clinical perspective.³ Clients' own behavioural and environmental factors also play a role in making care more complex in the presence of comorbidities.³ As Figure 1 neatly illustrates, all these factors contribute to the clinical complexity of treating comorbid chronic conditions, and thus impact the quality-of-care clients receive.³

Understanding how diseases cluster together is important for designing appropriate clinical disease management guidelines for certain chronic conditions that tend to co-occur.⁷ Certain chronic conditions tend to cluster together and which conditions cluster together may vary by gender. This is depicted in Figure 2 where Schäfer et al. (2014) used German insurance claims data to analyse which chronic conditions tend to be associated with one another on an individual level and of these diseases with high associations, which tend to be present together as multimorbidities.⁷



Figure 2: Disease associations in multimorbidity clusters for female population⁷




Figure 2 shows that, in the population considered, hypertension, lipid metabolism disorders, and chronic lower back pain had the highest number of associations with other chronic conditions among women.⁷ Figure 2 also shows the multitudes of associations between diseases classified as metabolic syndrome (such as obesity, hypertension, lipid metabolism disorders, and diabetes) and cardiovascular diseases (such as hypertension, atherosclerosis, chronic ischemic heart disease).⁷ Using this diagram Schäfer et al. (2014) clearly visually illustrate the complexity of how chronic conditions are associated and the multitude of combinations in which they can co-occur.⁷

The complexities of managing patients with comorbidities are evident in the higher rate of hospitalisation among medical scheme members who are registered for more than one NCD. Table 1 shows the risk of having been hospitalised in the previous 6-months among members with NCDs as per medical scheme chronic benefit registration data^b, compared to members who are not registered for an NCD but are registered for at least one chronic benefit. The results in Table 1 show that as the number of NCDs a medical scheme member is registered for increases, so does the risk of having been hospitalised in the previous six months. Members registered with one NCD are at 2.73 times higher risk of having been hospitalised compared to members who are not registered for any NCDs. This risk increases to 12.33 times for those registered for five or more chronic conditions.

^b NCDs included are diabetes, hypertension, heart attacks, coronary disease, mental disorders, asthma and arthritis.



Table 1: Risk ratio for hospitalisation in previous 6-months, compared to beneficiaries registered for chronic benefits other than NCDs (Own Analysis, Medical Scheme Data)

Number of NCDs	Risk ratio for having been hospitalised in the previous 6-months
1	2.73
2	4.01
3	6.07
4	8.30
5 +	12.33

The results in Table 1 are a clear illustration of how multimorbidity drives up the cost of care, and this increased risk of hospitalisation is also likely an indication of the increased complexity of caring for patients with multiple different chronic illnesses. This complexity is also driven by the fact that clinical management guidelines are not always explicit about how patients with comorbidities should be treated, given that medications for different chronic conditions may have varying levels of efficacy or side effects when taken in combination with one another.⁸

Polypharmacy, which is the regular intake of five or more medications, is often the outcome of the clinical complexity of managing multiple chronic conditions and although it is often appropriate or required, it also has additional risks particularly for the elderly, frail, or cognitively impaired.⁸ For instance, elderly patients taking more than four medications are at higher risk of falls with each additional medication, regardless of the medication type.⁹ For patients, polypharmacy is associated with reduced quality of life, increased risk of problems with mobility, increased contact with and dependence on the healthcare system, and higher risk of medication non-adherence, disability, adverse drug events, inappropriate use of medication, and long-term care placement.⁹ From a quality of care perspective, polypharmacy increases clinician workload and reduces their productivity, and also increases the likelihood of medication errors.⁹ Overall, polypharmacy increases the cost and burden to the healthcare system.⁹

When a clinician is not actively managing this load and assisting the client with ways to simplify the mental burden, adherence is likely to suffer- increasing the severity of some or all the conditions (Vignette 1).

Vignette 1: Medication overwhelm

The medication overwhelm for patients with co-morbid NCDs was illustrated most powerfully in one consultation I observed – between a dietician (D) and a woman in her late fifties (W). The woman arrived at the clinic wearing slippers and a grey hoodie. She removed both to be weighed and then took a seat in front of the dietician.

D: *Why have you come to see me today?*

W: *It's my kidneys and my heart. I was short of breath. So, I went to see the doctor and found out that my heart and my kidneys aren't working. They said I must come speak with you about my diet.*

The patient paused for a moment and then pulled out a shopping packet, full of medication:

W: *I don't know which pills are for my kidneys and which are for my heart. There's such a pile of pills, I don't know what's what.*

D: *'Bring them, I'll try.* [She looked closely at the packets, boxes, and blister packs] *'Okay, this one is for cholesterol, take it in the evening.'*



W: [The woman nodded] *In the evening.*

D: *Do you know what cholesterol is?*

W: [The woman shook her head] *No.*

D: *We'll speak about that later. What about this pill? Do you know it? This one is for blood pressure.*

W: *Okay, so I know the ones in the boxes, but I don't know the ones in these packets.'*

D: [The dietician scanned the labels] *Okay, this one is for heartburn [holding up a packet]. Can I write that on the label? Take it in the evening... Okay, next one: this one is also for high blood pressure. You take it at lunchtime. So, you have two...no three, three for blood pressure.*

By now, the dietician had stopped telling the patient when to take each pill. Even she was becoming overwhelmed. Nevertheless, she kept going:

D: *Okay, this is iron. That could be because your kidneys don't work so well, so you have too little iron.'*

W: [Now, it was the patient's turn to point at a box] *And this one, I know this one is for the heart.*

D: *Okay. And where's your water pill? Oh, here it is. I'll write it on the package.... And this one, I think if I had to guess, this one is for the heart that doesn't work very well. Yes, this is for the heart. [Slowly, the dietician returned the medicines back to the shopping packet] It's good to ask, the nurses at the pharmacy must explain to you. It's important that you ask. It's your body.*

But even she seemed to acknowledge that what was being asked of her patient – a caregiver of three grandchildren – was near impossible.

A study of patient and healthcare provider experiences with the care and management of comorbid chronic diseases at public healthcare facilities providing HIV care across Cape Town found that people living with HIV (PLWHIV) stated that they were inconvenienced by having to collect ART and other chronic medications at different facilities.¹⁰ They also complained of a lack of continuity of care as they were treated by different clinicians at every visit.¹⁰ Healthcare providers also stated that patients often struggled with the high pill burden.¹⁰

The risk that comes with multiple medications is lessened when someone is in the care of a team of clinicians who are aware of that individual's clinical history. In the private sector, it is possible for someone to have a designated general practitioner (GP) who supports them. However, GPs mostly work alone, and data is not shared freely between clinicians if a person needs to change GPs or expand her care to a broader care team. In the public sector, while healthcare workers demonstrate better teamwork, a client cannot 'book' with a particular health care worker. Therefore, clinicians are reliant on the robustness and depth of clinical notes left by other clinicians, usually documented in paper-based folders. This method of documentation makes patient record keeping more cumbersome and notes are often less comprehensive than required for streamlined care. Therefore, both the public and private systems currently do not lend themselves to quality care for either chronic or comorbid clients. Patients experience of care and the absolute level of quality of care provided would improve if the South African health system (private and public sectors) could roll out a single electronic health record.

The prevalence of comorbidities is difficult to measure

Data on chronic diseases (except HIV) is scarce in the South African public health sector context. The public health sector does not have unique patient identifiers which makes it impossible to know the true prevalence of individual chronic conditions and how these conditions interact in people and places. Routine administrative data systems currently only collect data on screenings conducted, with no collection of how many of the screens result in positive case finding. We therefore rely on survey data to estimate the prevalence of co- and multimorbidity of chronic conditions.



Public sector population

Without public sector data, we need to look to South African Demographic Health Survey (SADHS) of 2016 to assess comorbidity in the general population. Only four chronic illnesses were included in the results presented in Figure 3: diabetes, hypertension (high blood pressure), HIV, and cancer. The 2016 SADHS uses objective measures of HbA1C, blood pressure, and HIV tests to ascertain the prevalence of diabetes, hypertension (high blood pressure), and HIV status, respectively, while cancer is self-reported.¹¹ Respondents who stated that they were receiving cancer treatment or had been treated for cancer in the past are included in the results presented.

Although objective measures are obtained through tests, they are voluntary, therefore they do not necessarily represent the population prevalence of these chronic conditions, but rather only the prevalence among the population who volunteered to be tested. People may have refused to be tested because they perceived themselves to be in good health, or they may have feared disclosing their health status. In addition to this, only four chronic conditions are included in this analysis, many others have no objective measures in this dataset therefore they have not been included. These results, therefore, should not be taken to be a true representation of the prevalence of these diseases or multimorbidity, as they are likely to be underestimated.

Figure 3 shows that 57% of respondents were found to have no chronic conditions, 33% had at least one, 10% had two, while only 1% had three chronic conditions. Women had a higher prevalence of comorbidities than men, with 13% having two or three chronic illnesses, while 9% of men in the sample were found to have two or three chronic illnesses. Only one (female) respondent in the SADHS dataset had all four chronic conditions. Of all the respondents that had chronic conditions, 22.3% had one other comorbidity, and 1.6% had two or more other comorbidities.

Figure 3: Prevalence of comorbidity, SADHS 2016¹¹





Figure 4 shows the share of SADHS respondents with either diabetes, cancer, HIV, or hypertension that have at least one other chronic illness. The graph shows that 76% of diabetics, 57% of people who are being treated for cancer or have had cancer in the past, 46% of people living with HIV, and 29% of people with hypertension have at least one other chronic illness.



Figure 4: Proportion of respondents with a chronic illness that have at least one other chronic illness, SADHS 2016¹¹



Although survey data helps us to quantify the prevalence of the comorbidity of chronic illnesses in South Africa, Vignette 2 more vividly illustrates the lived experience of chronicity and comorbidity in one of South Africa's small, rural towns.

South Africa also routinely publishes statistics on causes of deaths. Unfortunately, cause of death data is not a good indicator of the way people lived. Although we have seen a shift in the cause of mortality from infectious to chronic diseases over time, the impact of NCDs is still likely understated in the cause of death data.



Vignette 2: Comorbidity on Farms

In December of 2019, I spent a day following the mobile clinic as it travelled from farm to farm, delivering medication to workers. Plumes of dust billowed out from behind the boxy white van as we travelled the twenty kilometres of dirt road to reach our first stop. We parked the van in front of a row of mud-brown cottages, in a rare strip of tree shade.

Medicines were wrapped in brown paper bags, passed from the back of the van like lunch packs. Those on insulin were handed bundles of plastic syringes with orange pistons. Some patients were weighed; while others had their blood pressure read, using a hand-pumped monitor and a well-placed stethoscope. On that farm alone, the list of patients was as follows:

Single NCD	Comorbid NCDs	Comorbid NCD and HIV/TB	Only HIV
Sonya, a middle-aged woman, collecting high blood pressure medication and requesting pain killers.	Annette, another middle-aged woman in a headscarf, living with co-morbid hypertension , diabetes and schizophrenia . She too would request pain medication.	Rita, a domestic worker, who emerged from the farmer's residence, and was handed HIV and hypertension medication.	Andre, a middle-aged man on HIV -treatment, who complained of lower back and chest pain .
Thembeke and Luthando, a husband and wife, both with hypertension , and requesting pain killers.	Nandi, a young woman (perhaps in her thirties) being treated for bipolar and hypertension .	Ettie, a young domestic worker who had recently recovered from TB , was in remission from leukemia , and managing hypertension .	
Lappies, a man in his forties, collecting high blood pressure medication			
Miriam, a diabetic domestic worker on insulin.			

We closed up the van and drove on to the next farm. Over the course of that day, we would see another twenty-or-so farm workers with much the same profile of cases. There was not a farm we visited that chronic illness had not touched. Not one without diabetes, and nearly everyone with hypertension. Always a mix of infectious and non-infectious illnesses – sometimes in the same households, even occupying the same bodies

Medical scheme population

In the private health sector, unlike the public sector, unique patient identifiers are available. However, the industry is fragmented and reliant on the Council for Medical Schemes (CMS) to collate and report on medical scheme population data. The CMS produces annual statistics that track prevalence of chronic conditions, including comorbidity, and this is helpful in understanding the trend of chronic disease in this population.




Using the CMS data, we have counted the number of people with at least one claim for two or more chronic illness (this includes everyone who claimed for a chronic condition at least once in the past year) to estimate the number of medical scheme members with comorbidities. It should be noted that this estimate is likely to underestimate the true prevalence of comorbidities in this population for two reasons. Firstly, not all medical aid members will have been screened for chronic illnesses, so they may be unaware of their conditions. Secondly, not all medical aid members will claim from their medical schemes when purchasing chronic medication or care for chronic illnesses, many will simply pay out of pocket.

The medical scheme population is older than the general South African population and also differs in terms of socioeconomic status. As we have explained in earlier briefs within this series, both age and socioeconomic status have an impact on the prevalence, incidence, and severity of NCDs. Although older people are more likely to suffer from chronic conditions, people of higher socioeconomic status are more likely to have more education, access to better nutrition and better-quality healthcare and thus tend to be in better health. Therefore, it is difficult to say for sure whether the prevalence of comorbidities is higher or lower in the medical scheme population compared to the general population. However, the prevalence of HIV is much higher in the non-medical scheme population, therefore it is likely that the prevalence of comorbidities related to HIV is higher in this population.

Recently, the CMS has begun to publish a report on chronic conditions and co-occurrence with one another, which has shed light on the prevalence of comorbidity in the insured population. Figure 5 shows the rates of two, three and four or more comorbidities in the medical scheme environment. The data shows an increasing trend between 2011 and 2017 for all numbers of comorbidities. While still being the least prevalent, individuals with four or more chronic conditions has increased by 91% since 2011 (0.22% to 0.42% in 2017). Individuals with three or more chronic conditions has shown a 50% increase between 2011 and 2017- a substantial jump in just six years. The medical scheme population shows an overall prevalence of 6% of people (~556,160 individuals) living with comorbid chronic conditions. This is lower than the 11% estimated for the general population using the SADHS survey data.

Figure 5: Comorbidity burden in medical scheme population¹²



Overall, 44% of members who were registered for a chronic condition were registered for two or more chronic conditions. Very worryingly, 85% of medical scheme members with diabetes also had one or more other chronic conditions (see Table 2).



Table 2: Co-morbidity according to chronic registration for NCDs in medical schemes (Own Analysis, Medical Scheme Data)

NCD	Percentage of those with the NCD that have 1 or more other NCDs
Diabetes	85%
Hypertension	55%
Heart attack	92%
Coronary disease	89%
Mental disorders	66%
Asthma	53%
Arthritis	85%
Any NCD	44%

From our mental health brief in this series, we showed how chronic conditions co-occur with mental health conditions. Mental health conditions can make it difficult for people to adhere to their other chronic medication (and even their mental health related medication). This makes mental health a key predictor of health outcomes for people living with chronic conditions. Furthermore, mental health conditions, particularly depression and anxiety, often arise after diagnosis with diabetes, hypertension, or HIV.^{13,14} Therefore, awareness and screening for mental health conditions in those with other chronic conditions should be a basic standard of care.

Using data obtained from one of South Africa's largest medical schemes, Table 3 shows the risk ratios, by sex, of having a mental health condition given that one already has one of the listed chronic conditions. A risk ratio greater than one indicates an increased risk. Therefore, the table shows that people with asthma, heart problems, hypertension, diabetes, and arthritis are at higher risk of having comorbid mental health conditions relative to those without these chronic illnesses. Compared to the other chronic illnesses, people with arthritis are at highest risk of having comorbid mental health conditions, as they are 3.94 times as likely to have mental health conditions compared to those without arthritis. Men are also at higher risk of having comorbid mental health conditions for every chronic illness, except for diabetes, when compared to women, but perhaps most notable is the increased risk for all five of the conditions in the table.

Table 3: Risk ratios for people living with a mental health condition (own analysis using medical scheme data)^c

	Asthma	Heart Problems	Hypertension	Diabetes	Arthritis
Female	2,62	2,55	1,83	1,78	2,91
Male	2,77	2,91	1,90	1,73	3,94
Both	2,76	2,49	1,84	1,69	3,42

A person-centred approach is required

^c Data obtained from one of South Africa's largest medical schemes



Managing the care of people with co- or multimorbidities requires a whole person-centred approach as opposed to simply treating the disease. As illustrated in Figure 1, there are not only clinical considerations but also behavioural and environmental factors which need to be accounted for when managing multiple chronic conditions.

Figure 2 also shows the relationship between chronic illnesses and the patterns of how these cluster together. Clinical guidelines need to take this into account so that more holistic disease management approaches are designed.⁸ To do this, more integrated care approaches need to be adopted. These approaches should allow patients to receive treatment in ways which enable continuity of care and track which medications have been prescribed to minimise polypharmacy where possible. Tools such as STOPP (screening tool of older people's prescriptions) and START (screening tool to alert to right treatment) criteria can be used to assist with decision making.⁹ These tools help clinicians to compare medication lists so that duplication can be identified, and have been found to significantly reduce inappropriate prescriptions.⁹ They are also useful for tailoring treatment to the needs of specific individuals and prevent resource wastage caused by repeat prescriptions that are not regularly re-evaluated.


Looking to the future

For a country to understand its comorbidity burden, it needs to be able to track individuals in the health system through a unique patient identifier or electronic health record (EHR). This would allow the health system to know whether someone has been screened, what the result was and whether they are on a medication/treatment regimen as a result. As more conditions arise, an EHR would allow the health system to alert health care workers that the person is a high-risk individual requiring more intensive care. As mentioned, only South Africa's medical scheme industry has this system in place.

As part of this work, Percept plans to forecast the burden of chronic disease between 2021-2030. The scarcity of accurate data in the South Africa health system makes this modelling exercise all the more difficult. With the onset of Covid-19, we have seen how important accurate, individual-level patient data is for being able to manage the pandemic. Covid-19 has far worse outcomes for those with certain chronic conditions, and even more so for those with multiple chronic conditions and therefore it would be best for this subset of the population to be vaccinated first. However, we simply do not have the data available to do so.

Furthermore, most often, disease modelling is focused on a single disease, ignoring the interactions between diseases themselves and how these may influence growth rates. The interconnectedness of chronic diseases means that one cannot ignore these influences if you want to accurately estimate the current and future burden of chronic disease.

Figure 6: Resource use relative to clinical complexity^d



In this brief we explained 'super-additivity', which is where the sum of multiple conditions costs more than the parts, and how chronic conditions are often associated with each other and cluster together. Figure 6 illustrates that resource use is most intense for the small group of people who are the most clinically complex. This is very often those with comorbidities (chronic or otherwise). This means that once we have been able to forecast the most prevalent chronic diseases, we will also be able to determine which conditions should be targeted for prevention activities, to generate the best health outcomes at the lowest possible cost. This can help to free up resources for primary health care, which helps to keep the

^d Figure designed by PPO Serve. Ruff, B. 2018



population well.

Conclusion

In this brief we have highlighted both the interconnectedness of different chronic conditions and the care that these complex clients require. To provide quality care, the South African health systems need to adapt, allowing for better record keeping, easier sharing of data and comprehensive clinical care. Certain chronic conditions are predictors of others and therefore, by targeting one or two of the most prevalent conditions, we could bring down the number of comorbidities at the individual and collective level.

References

1. Valderas J, Starfield B, Sibbald B, Salisbury C, Roland M. Defining Comorbidity: Implications for Understanding Health and Health Services. *Ann Fam Med*. 2009;7(4):357-363. doi:10.1370/afm.983
2. Biswas T, Townsend N, Islam MS, et al. Association between socioeconomic status and prevalence of non-communicable diseases risk factors and comorbidities in Bangladesh: findings from a nationwide cross-sectional survey. *BMJ Open*. 2019;9(9). doi:10.1136/bmjopen-2018-025538
3. Zulman DM, Asch SM, Martins SB, Kerr EA, Hoffman BB, Goldstein MK. Quality of care for patients with multiple chronic conditions: The role of comorbidity interrelatedness. *J Gen Intern Med*. 2014;29(3):529-537. doi:10.1007/s11606-013-2616-9
4. Gouda HN, Charlson F, Sorsdahl K, et al. Burden of non-communicable diseases in sub-Saharan Africa: results from the Global Burden of Disease Study 2017. *Lancet Glob Heal*. 2019;7(10):E1375-E1387. doi:10.1016/S2214-109X(19)30374-2
5. Cortaredona S, Ventelou B. The extra cost of comorbidity: Multiple illnesses and the economic burden of non-communicable diseases. *BMC Med*. 2017;15(1):1-11. doi:10.1186/s12916-017-0978-2
6. Blakely T, Kvizhinadze G, Atkinson J, Dieleman J, Clarke P. Health system costs for individual and comorbid noncommunicable diseases: An analysis of publicly funded health events from New Zealand. *PLoS Med*. 2019;16(1):e1002716. doi:10.1371/journal.pmed.1002716
7. Schäfer I, Kaduszkiewicz H, Wagner H-O, Schön G, Scherer M, Van Den Bussche H. *Reducing Complexity: A Visualisation of Multimorbidity by Combining Disease Clusters and Triads*; 2014. doi:10.1186/1471-2458-14-1285
8. Guthrie B, Payne K, Alderson P, McMurdo ME, Mercer S. Adapting clinical guidelines to take account of multimorbidity. *Br Med J*. 345AD;4:e6341-e6341. Accessed May 24, 2021. <https://eprints.gla.ac.uk/70694/1/70694.pdf>
9. Halli-Tierney A, Scarbrough C, Carroll D. Polypharmacy: Evaluating Risks and Deprescribing. *Am Fam Physician*. 2019;100(1):32-38. Accessed May 24, 2021. <https://www.aafp.org/afp/2019/0701/p32.html>
10. Id NP, De Villiers A, Jonathan D, Kalombo C, Kengneid A-P. Care and management of a double burden of chronic diseases: Experiences of patients and perceptions of their healthcare providers. Published online 2020. doi:10.1371/journal.pone.0235710
11. National Department of Health (NDOH), Statistics South Africa (Stats SA), South African Medical Research Council (SAMRC), ICF. *South Africa Demographic and Health Survey 2016*; 2019.
12. Council for Medical Schemes. Prevalence of chronic diseases in the population covered by medical aid schemes in South Africa. 2019;(March).
13. Madavanakadu Devassy S, Madavanakadu Devassy S, Benny AM, et al. Social factors associated with chronic non-communicable disease and comorbidity with mental health problems in India: A scoping review. *BMJ Open*. 2020;10(6):e035590. doi:10.1136/bmjopen-2019-035590



14. Nel A, Kagee A. The relationship between depression, anxiety and medication adherence among patients receiving antiretroviral treatment in South Africa. *AIDS Care - Psychol Socio-Medical Asp AIDS/HIV*. 2013;25(8):948-955. doi:10.1080/09540121.2012.748867